

DESY 02-220
 hep-ph/0301177

QCD-INSTANTONS AND SATURATION AT SMALL x^*

F. SCHREMPP AND A. UTERMANN

*Deutsches Elektronen Synchrotron (DESY),
 Notkestrasse 85,
 D-22607 Hamburg, Germany*

E-mail: fridger.schrempp@desy.de, andre.utermann@desy.de

We briefly report on the contribution of QCD-instantons to the phenomenon of saturation in deep-inelastic scattering (DIS) at small Bjorken- x . The explicitly known instanton gauge field serves as a concrete realization of an underlying non-perturbative saturation mechanism associated with strong classical fields. Two independent strategies were pursued, with consistent results: On the one hand, an approach starting from instanton-perturbation theory was considered and on the other hand, the scattering of a Wilson loop in an instanton gauge field background. In both cases, the conspicuous, intrinsic instanton size scale $\langle \rho \rangle \approx 0.5$ fm, as known from the lattice, turns out to determine the saturation scale.

1. Setting the stage

eP -scattering at small Bjorken- x uncovers a novel regime of QCD, where the coupling α_s is small, but the parton densities are so large that conventional perturbation theory ceases to be applicable. While there is experimental evidence from HERA for a strong growth of the gluon density at small x_{Bj} , a mechanism is needed that eventually leads to “saturation”, i.e. to a limitation of the maximum gluon density per unit of phase space. Much interest has recently been generated through association of the saturation phenomenon with a multiparticle quantum state of high occupation numbers, the “Colour Glass Condensate” that correspondingly, can be viewed as a strong *classical* colour field ¹.

In this paper, we briefly report on the promising rôle of QCD-instantons ² (I) in this context. With the help of crucial information from lattice simulations ^{3,4}, we consider a background instanton as an explicitly known, truly non-perturbative classical gauge field ($\propto 1/g_s$) in the context of saturation at small x_{Bj} . A crucial aspect concerns the I -size ρ . On the one hand it is just a collective coordinate to be integrated over

*Talk presented at the Workshop on Strong and Electroweak Matter (SEWM 2002), October 2-5, 2002, Heidelberg, Germany.

in any observable, with the I -size distribution $D(\rho) = dn_I/d^4z d\rho$ as universal weight. On the other hand, according to lattice data, $D(\rho)$ turns out to be *sharply* peaked (Fig. 1 (left)) around $\langle\rho\rangle \approx 0.5$ fm. Hence instantons represent truly non-perturbative gluons that bring in naturally an intrinsic size scale $\langle\rho\rangle$ of hadronic dimension. As we shall see, $\langle\rho\rangle$ actually determines the saturation scale ^{6,7}. Presumably, it is also reflected in the conspicuous *geometrization* of soft QCD at high energies ^{8,6,7}. For related approaches associating instantons with high-energy (diffractive) scattering, see Refs. ^{9,10,11,12}.

We know already from I -perturbation theory that the instanton contribution tends to strongly increase towards the soft regime ^{13,14,15}. The mechanism for the decreasing instanton suppression with increasing energy is known since a long time ^{16,12}: Feeding increasing energy into the scattering process makes the picture shift from one of tunneling between vacua ($E \approx 0$) to that of the actual creation of the sphaleron-like, coherent multi-gluon configuration ¹⁷ on top of the potential barrier of height ¹³ $E = M_{\text{sph}} \propto \frac{1}{\alpha_s \rho_{\text{eff}}}$. In Ref.⁷, we have already argued by means of lattice results that the QCD - “sphaleron” is playing an essential rôle in building up the “Colour Glass Condensate”.




Figure 1. (Left) UKQCD lattice data^{3,4,5} of the $(I + \bar{I})$ -size distribution for quenched QCD ($n_f = 0$). Both the sharply defined I -size scale $\langle\rho\rangle \approx 0.5$ fm and the parameter-free agreement with I -perturbation theory^{4,5} (solid line) for $\rho \lesssim 0.35$ fm are apparent. (Right) the simplest I -induced process ²⁰ transcribed into the colour dipole framework.

The colour dipole picture¹⁸ represents an intuitive framework for investigating saturation effects at small x_{Bj} . In the proton's rest frame, the virtual photon fluctuates mainly into a $q\bar{q}$ -dipole a long distance upstream of the target proton. The large difference of the $\gamma^* \rightarrow q\bar{q}$ -dipole formation and $(q\bar{q})$ - P interaction times at small x_{Bj} generically give rise to the familiar factorized expression of the inclusive photon-proton cross sections,

$$\sigma_{L,T}(x_{\text{Bj}}, Q^2) = \int_0^1 dz \int d^2\mathbf{r} |\Psi_{L,T}(z, r)|^2 \sigma_D(r, \dots). \quad (1)$$

Here, $|\Psi_{L,T}(z, r)|^2$ denotes the modulus squared of the (light-cone) wave function of the virtual photon, calculable in pQCD, and $\sigma_D(r, \dots)$ is the $q\bar{q}$ -dipole-nucleon cross section. The variables in Eq. (1) are the transverse $(q\bar{q})$ -size $\mathbf{r}$ and the photon's longitudinal momentum fraction z carried by the quark. The dipole cross section is expected to include in general the main non-perturbative contributions. For small r , one finds within pQCD^{18,19} that σ_D vanishes with the area πr^2 of the $q\bar{q}$ -dipole. Besides this phenomenon of ‘‘colour transparency’’ for small r , the dipole cross section is expected to saturate towards a constant, once the $q\bar{q}$ -separation r exceeds a certain saturation scale r_s .

2. Instantons and Saturation

2.1. Starting from Instanton Perturbation Theory

Next, let us briefly illustrate that in the presence of a background instanton, the saturation scale r_s indeed equals the intrinsic I -size scale $\langle \rho \rangle$. For reasons of space, we restrict the discussion to the simplest I -induced process²⁰, $\gamma^* g \Rightarrow q_R \bar{q}_R$, with one flavour and no final-state gluons (Fig. 1 (right)). More details and the realistic case with gluons and three light flavours, using the $I\bar{I}$ -valley approach, may be found in Ref.⁷. The strategy is to transform the known results on I -induced processes in DIS into the colour dipole picture. By exploiting the lattice results on the I -size distribution (Fig. 1 (left)), we then carefully increase the $q\bar{q}$ -dipole size r towards hadronic dimensions. Let us start by recalling the results from Ref.²⁰,

$$\sigma_{L,T}(x_{\text{Bj}}, Q^2) = \int_{x_{\text{Bj}}}^1 \frac{dx}{x} \left(\frac{x_{\text{Bj}}}{x} \right) G\left(\frac{x_{\text{Bj}}}{x}, \mu^2\right) \int dt \frac{d\hat{\sigma}_{L,T}^{\gamma^*g}(x, t, Q^2)}{dt}; \quad (2)$$

$$\frac{d\hat{\sigma}_L^{\gamma^*g}}{dt} = \frac{\pi^7}{2} \frac{e_q^2}{Q^2} \frac{\alpha_{\text{em}}}{\alpha_s} \left[x(1-x)\sqrt{tu} \frac{\mathcal{R}(\sqrt{-t}) - \mathcal{R}(Q)}{t + Q^2} - (t \leftrightarrow u) \right]^2 \quad (3)$$

and a similar expression for $d\hat{\sigma}_T^{\gamma^*g}/dt$.

Eqs. (2, 3) involve the master integral $\mathcal{R}(\mathcal{Q})$ with dimensions of a length,

$$\mathcal{R}(\mathcal{Q}) = \int_0^\infty d\rho D(\rho) \rho^5 (\mathcal{Q}\rho) K_1(\mathcal{Q}\rho). \quad (4)$$

By means of an appropriate change of variables and a subsequent $2d$ -Fourier transformation, Eqs. (2, 3) may indeed be cast ⁷ into a colour dipole form (1), e.g. (with $\hat{Q} = \sqrt{z(1-z)}Q$)

$$\begin{aligned} \left(|\Psi_L|^2 \sigma_D\right)^{(I)} &\approx |\Psi_L^{\text{pQCD}}(z, r)|^2 \frac{1}{\alpha_s} x_{\text{Bj}} G(x_{\text{Bj}}, \mu^2) \frac{\pi^8}{12} \\ &\times \left\{ \int_0^\infty d\rho D(\rho) \rho^5 \left(\frac{-\frac{d}{dr^2} \left(2r^2 \frac{K_1(\hat{Q}\sqrt{r^2+\rho^2/z})}{\hat{Q}\sqrt{r^2+\rho^2/z}} \right)}{K_0(\hat{Q}r)} - (z \leftrightarrow 1-z) \right) \right\}^2. \end{aligned} \quad (5)$$

The strong peaking of $D_{\text{lattice}}(\rho)$ around $\rho \approx \langle \rho \rangle$, implies

$$\left(|\Psi_{L,T}|^2 \sigma_D\right)^{(I)} \Rightarrow \begin{cases} \mathcal{O}(1) \text{ but exponentially small;} & r \rightarrow 0, \\ |\Psi_{L,T}^{\text{pQCD}}|^2 \frac{1}{\alpha_s} x_{\text{Bj}} G(x_{\text{Bj}}, \mu^2) \frac{\pi^8}{12} \mathcal{R}(0)^2; & \frac{r}{\langle \rho \rangle} \gtrsim 1. \end{cases} \quad (6)$$

Hence, the association of the intrinsic instanton scale $\langle \rho \rangle$ with the saturation scale r_s becomes apparent from Eqs. (5, 6): $\sigma_D^{(I)}(r, \dots)$ rises strongly as function of r around $r_s \approx \langle \rho \rangle$, and indeed *saturates* for $r/\langle \rho \rangle > 1$ towards a *constant geometrical limit*, proportional to the area $\pi \mathcal{R}(0)^2 = \pi \left(\int_0^\infty d\rho D_{\text{lattice}}(\rho) \rho^5 \right)^2$, subtended by the instanton. Since $\mathcal{R}(0)$ would be divergent within I -perturbation theory, the information about $D(\rho)$ from the lattice (Fig. 1 (left)) is crucial for the finiteness of the result.

2.2. Wilson-Loop Scattering in an Instanton Background

Complementary to the above strategy of extending the known results of I -perturbation theory by means of lattice information into the saturation regime, we have followed a second promising route ²¹. As a simple semi-classical eikonal estimate of the total $q\bar{q}$ -dipole cross section, the scattering of a pair of infinitely long Wilson lines ($q\bar{q}$ -dipole) on the proton in an instanton background field was evaluated along the lines of Refs. ^{22,10,11},

$$\sigma_D^{(I)}(r) = \int d^2\mathbf{b} \text{Im} T(\mathbf{b}, \mathbf{r}) = \int d^2\mathbf{b} \frac{2}{N_c} \text{tr} \left\langle \text{Re} \{ (1 - S^{(I)}(\mathbf{b}, \mathbf{r})) T_P \} \right\rangle_{G^{(I)}} \quad (7)$$

The functional integration over the I -background field in Eq. (7) reduces to integrations over the I -collective coordinates,

$$\langle \dots \rangle_{G^{(I)}} : \mathcal{D}G_\mu^a \Rightarrow d^4z d\rho D_{\text{lattice}}(\rho) dU, \quad (8)$$

with z_μ denoting the I -position 4-vector and U the I -colour orientation. The Wilson loop in Eq. (7), associated with the $q\bar{q}$ -dipole,

$$S^{(I)}(\mathbf{b}, \mathbf{r}) = W^\dagger(\mathbf{b}) W(\mathbf{b} + \mathbf{r}), \quad (9)$$

$$W(\mathbf{x}_\perp) = P \exp \left\{ i g_s \int_{-\infty}^{\infty} d\tau v \cdot G^{(I)}(\tau v + x_\perp) \right\}; \quad \begin{aligned} v_\mu &= q_\mu/Q, \\ x_\perp \cdot v &= 0, \end{aligned} \quad (10)$$

turns out to only involve an integration over the z_+ light cone component of the I -position, while the remaining $dz_- d^2\mathbf{z}$ integrations only act on the factor $T_P(\dots)$, associated with the proton structure. Again, the result can be obtained in analytic form with similar features as in our previously discussed approach based on I -perturbation theory. The predicted ratio $\sigma_D^{(I)}(r)/\sigma_D^{(I)}(\infty)$ displayed in Fig. 2 (left) as function of $r/\langle\rho\rangle$, illustrates the importance of $\langle\rho\rangle$ in the approach to saturation. In Fig. 2 (right), the corresponding impact parameter profile for $r = \langle\rho\rangle, \infty$ is shown. The important issue of understanding the x_{Bj} -dependence of the saturation scale within the present I -approach, is currently under active investigation ²¹.




Figure 2. (Left) approach to saturation around $r \gtrsim \langle\rho\rangle$ and (right) central b -profile

Acknowledgements

We thank Igor Cherednikov and Michael Lublinsky for interesting discussions.

References

1. E. Iancu, A. Leonidov and L. D. McLerran, *Nucl. Phys.* **A692**, 583 (2001).
2. A. Belavin *et al.*, *Phys. Lett.* **B59**, 85 (1975).
3. D.A. Smith and M.J. Teper (UKQCD), *Phys. Rev.* **D58**, 014505 (1998).
4. A. Ringwald and F. Schrempp, *Phys. Lett.* **B459**, 249 (1999).
5. A. Ringwald and F. Schrempp, *Phys. Lett.* **B503**, 331 (2001).
6. F. Schrempp and A. Utermann, *Acta Phys. Polon.* **B33**, 3633 (2002).
7. F. Schrempp and A. Utermann, *Phys. Lett.* **B543**, 197 (2002).
8. F. Schrempp, *J. Phys.* **G28**, 915 (2002).
9. D.E. Kharzeev, Y.V. Kovchegov and E. Levin, *Nucl. Phys.* **A690**, 621 (2001).
10. E. Shuryak and I. Zahed, *Phys. Rev.* **D62**, 085014 (2000).
11. M. A. Nowak, E. V. Shuryak and I. Zahed, *Phys. Rev.* **D64**, 034008 (2001).
12. D.M. Ostrovsky *et al.*, *Phys. Rev.* **D66**, 036004 (2002).
13. A. Ringwald and F. Schrempp, *Proc. Quarks '94*, ed D.Yu. Grigoriev *et al.* (Singapore: World Scientific) p 170, [arXiv:hep-ph/9411217].
14. A. Ringwald and F. Schrempp, *Phys. Lett.* **B438**, 217 (1998).
15. A. Ringwald and F. Schrempp, *Comput. Phys. Commun.* **132**, 267 (2000).
16. H. Aoyama and H. Goldberg, *Phys. Lett.* **B188**, 506 (1987); A. Ringwald, *Nucl. Phys.* **B330**, 1 (1990); O. Espinosa, *Nucl. Phys.* **B343**, 310 (1990).
17. F. R. Klinkhamer and N. S. Manton, *Phys. Rev.* **D30**, 2212 (1984).
18. N. Nikolaev and B.G. Zakharov, *Z. Phys.* **C49**, 607 (1990); *Z. Phys.* **C53**, 331 (1992); A.H. Mueller, *Nucl. Phys.* **B 415**, 373 (1994).
19. F. E. Low, *Phys. Rev.* **D12**, 163 (1975);
L. Frankfurt, G.A. Miller and M. Strikman, *Phys. Lett.* **B304**, 1 (1993).
20. S. Moch, A. Ringwald and F. Schrempp, *Nucl. Phys.* **B507**, 134 (1997).
21. F. Schrempp and A. Utermann, in preparation.
22. A. Krämer and H. G. Dosch, *Phys. Lett.* **B252**, 669 (1990);
O. Nachtmann, *Annals Phys.* **209**, 436 (1991);
W. Buchmüller and A. Hebecker, *Nucl. Phys.* **B476**, 203 (1996);
A. I. Shoshi, F. D. Steffen and H. J. Pirner, *Nucl. Phys.* **A709**, 131 (2002).